

Download Study Guide Spencer Rathus

In human sexuality, the refractory period is usually the recovery phase after orgasm during which it is physiologically impossible for a man to have additional orgasms. This phase begins immediately after ejaculation and lasts until the excitement phase of the human sexual response cycle begins anew with low level response. Although it is generally reported that women do not experience a ...Sexual activity and innuendo. The pelvic thrust is used during sexual intercourse by many species of mammals, including humans, or for other sexual activities (such as non-penetrative sex).In 2007, German scientists noted that female monkeys could increase the vigor and amount of pelvic thrusts made by the male by shouting during intercourse. In whitetail deer, copulation consists of a single ...Cunnilingus is an oral sex act performed by a person on the female genitalia (the clitoris, other parts of the vulva or the vagina). The clitoris is the most sexually sensitive part of the human female genitalia, and its stimulation may result in female sexual arousal or orgasm.. Cunnilingus can be sexually arousing for participants and may be performed by a sexual partner as foreplay to ...Orgasm (from Greek ???????? orgasmos "excitement, swelling"; also sexual climax) is the sudden discharge of accumulated sexual excitement during the sexual response cycle, resulting in rhythmic muscular contractions in the pelvic region characterized by sexual pleasure. Experienced by males and females, orgasms are controlled by the involuntary or autonomic nervous system.